HEALTHY FOOD SYSTEMS

HEALTHY ENVIRONMENTS

### University of California

**Agriculture and Natural Resources** 



# UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

	WEEKLY SOIL MOISTURE LOSS IN INCHES												
	(Estimated Crop Evapotranspiration or ET <sub>C</sub> )												
04/11/25 through 04/17/25													
Crops (Leafout Date)	ŧ	148 Merce	d		#39 Parlier				#258 Lemon Cove				
	04/11 - 04/17	Accum'd	04/18 - 04/24		04/11 - 04/17	Accum'd	04/18 - 04/24		04/11 - 04/17	Accum'd	04/18 - 04/24	1	
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated	1	
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc	1	
Almonds (3/1) *	0.87	4.27	1.03		0.95	4.39	1.04		0.85	3.93	1.06	1	
Pistachio (TBD) * **	0.00	0.00	0.00		0.00	0.00	0.00		0.00	0.00	0.00	1	
Citrus (2/1)	0.80	6.13	0.90		0.89	6.39	0.91		0.79	5.81	0.93	1	
Raisin Grapes (4/14) (11 ft. row spacing)	0.03	0.03	0.08		0.03	0.03	0.08		0.02	0.02	0.08		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	0.07	0.07	0.14		0.06	0.06	0.14		0.05	0.05	0.15		
Walnuts (4/14)	0.08	0.08	0.14		0.08	0.08	0.14		0.07	0.07	0.14	1	
Stone Fruit (3/8)	0.45	1.59	0.55		0.49	1.74	0.55		0.43	1.57	0.57		
Past 7 days precipitation (inches)		0.00				0.00				0.00			
Accumulated precipitation (inches) (1/1/2025)		0.00				5.22				4.26		<u> </u>	

Dates in parentheses above, indicate leaf out or starting date for ET accumulation for the specific crop

\* Estimates are for orchard floor conditions where vegetation is managed by some combination of arrip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

\*\* Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.

\*\*\* Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

ty whe lear water potential reactes approximate -1.9 ML at Orowers should adjust the imagaton start date based on the individual vinteyard reaction and son type. BAST WEEKL V ADDI HED WATED IN INCHES AD HIGTED FOD EEEICIENCY

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY <sup>1</sup>												
Crops	#148 Merced					#39 Parlier			#258 Lemon Cove			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	1.3	1.2	1.0	0.9	1.5	1.3	1.1	1.0	1.3	1.1	1.0	0.9
Pistachio (TBD)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Citrus (2/1)	1.2	1.1	0.9	0.8	1.4	1.2	1.0	0.9	1.2	1.1	0.9	0.8
Raisin Grapes (4/14) (11 ft. row spacing)***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis) ***	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Walnuts (4/14)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Stone Fruit (3/8)	0.7	0.6	0.5	0.5	0.8	0.7	0.6	0.5	0.7	0.6	0.5	0.5

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE												
Crops		#148 Merced				#39 Parlier			#258 Lemon Cove			
Almonds 115 Trees/A	307	283	236	213	354	307	260	236	307	260	236	213
Pistachio 106 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0
Citrus 110 Trees/A	296	272	222	197	346	296	247	222	296	272	222	197
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0
Walnuts 76 Trees/A	36	36	36	36	36	36	36	36	36	36	36	36
Stonefruit 172 Trees/A	111	95	79	79	126	111	95	79	111	95	79	79
For further information concerning all counties receiving this report contac	t the Fresno (	Co. Farm Adv	isor's office a	t (559) 241-	7526							

#### HEALTHY FOOD SYSTEMS + HEALTHY ENVIRONMENTS + HEALTHY COMMUNITIES + HEALTHY CALIFORNIANS

## University of California

**Agriculture and Natural Resources** 

## UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

	WEEKLY SOIL MOISTURE LOSS IN INCHES												
	(Est	imated Cro	p Evapotrans	piration or 1	ET <sub>C</sub> )								
04/11/25 through 04/17/25													
Crops (Leafout Date) #124 Panoche #2 Five Points #15 Stratford													
	04/11-04/17	Accum'd	04/18- 04/24		04/11-04/17	Accum'd	04/18- 04/24		04/11-04/17	Accum'd	04/18- 04/24		
	Water	Seasonal	Estimated		Water	Seasonal	Estimated		Water	Seasonal	Estimated		
	Use	Water Use	ETc		Use	Water Use	ETc		Use	Water Use	ETc		
Almonds (3/1) *	1.03	4.06	1.16		1.04	4.23	1.21		1.11	4.51	1.21		
Pistachio (TBD) * **	0.00	0.00	0.00		0.00	0.00	0.00		0.00	0.00	0.00		
Citrus (2/1)	0.94	6.65	1.02		0.96	6.81	1.05		1.02	7.32	1.05		
Raisin Grapes (4/14) (11 ft. row spacing)	0.03	0.03	0.08		0.03	0.03	0.09		0.03	0.03	0.09		
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	0.07	0.07	0.16		0.06	0.06	0.17		0.07	0.07	0.18		
Walnuts (4/14)	0.09	0.09	0.18		0.09	0.09	0.17		0.09	0.09	0.21		
Stone Fruit (3/8)	0.53	1.78	0.63		0.53	1.72	0.65		0.58	1.80	0.65		
Past 7 days precipitation (inches)		0.00				0.00				0.00			
Accumulated precipitation (inches) (1/1/2025)		2.21				2.85				2.51			
Dates in parentheses above, indicate leaf out or starting date for ET accumulation t	or the specific cr	op		•					•				
* Estimates are for orchard floor conditions where vegetation is managed by some	combination of s	strip applicati	ons of herbicide	s, frequent m	owing or tillage	, and by mid a	and late season s	hading and w	vater stress. We	ekly estimates	s of soil moistur	e loss can be	
as much as 25 percent higher in orchards where cover crops are planted and manage													
** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.1	9 - resulting in a	bout 8% grea	ter water use tha	n shown in th	ese tables.								

\*\*\* Raisin Grapes and Winegrapes Irrigation should hold off until midday leaf water potential drops to -1.0 MPa, before that soil moisture reservoir is sufficient to supply the vine water demand. Update will be sent shortly once the county wide leaf water potential reaches approximate -1.0 MPa. Growers should adjust the irrigation start date based on the individual vineyard location and soil type.

 11	J	0	,	21
	PAST WEEKLY A	APPLIED WATER IN	INCHES ADJUST	ED FOR EFFICIENCY 1

Crops	#124 Panoche				5, ADJUST	#2 Five Poi			#15 Stratford			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%
Almonds (3/1)	1.6	1.4	1.2	1.1	1.6	1.4	1.2	1.1	1.7	1.5	1.3	1.2
Pistachio (TBD)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Citrus (2/1)	1.4	1.3	1.1	1.0	1.5	1.3	1.1	1.0	1.6	1.4	1.2	1.1
Raisin Grapes (4/14) (11 ft. row spacing)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Winegrapes (4/14) (10 ft. spacing on California Sprawl Trellis)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Walnuts (4/14)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Stone Fruit (3/8)	0.8	0.7	0.6	0.6	0.8	0.7	0.6	0.6	0.9	0.8	0.7	0.6

1 The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

	PAST WEEKLY APPLIED WATER IN GALLON PER TREE OR VINE												
Crops	#124 Panoche				#2 Five Points								
Almonds 115 Trees/A	378	331	283	260	378	331	283	260	401	354	307	283	
Pistachio 106 Trees/A	0	0	0	0	0	0	0	0	0	0	0	0	
Citrus 110 Trees/A	346	321	272	247	370	321	272	247	395	346	296	272	
Raisin Grapes 566 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0	
Winegrapes 622 Vines/A	0	0	0	0	0	0	0	0	0	0	0	0	
Walnuts 76 Trees/A	36	36	36	36	36	36	36	36	36	36	36	36	
Stonefruit 172 Trees/A	126	111	95	95	126	111	95	95	142	126	111	95	
For further information concerning all counties receiving this report, contact	the Fresno Co	o. Farm Advis	sor's office at	(559) 241-7	526.								